

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

MARK SCHEME for the October/November 2015 series**0625 PHYSICS****0625/32**

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2015 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0625	32

NOTES ABOUT MARK SCHEME SYMBOLS & OTHER MATTERS

- B marks** are independent marks, which do not depend on other marks. For a B mark to be scored, the point to which it refers must be seen specifically in the candidate's answer.
- M marks** are method marks upon which accuracy marks (A marks) later depend. For an M mark to be scored, the point to which it refers **must** be seen in a candidate's answer. If a candidate fails to score a particular M mark, then none of the dependent A marks can be scored.
- C marks** are compensatory marks in general applicable to numerical questions. These can be scored even if the point to which they refer are not written down by the candidate, **provided subsequent working gives evidence that they must have known it**. For example, if an equation carries a C mark and the candidate does not write down the actual equation but does correct substitution or working which shows he knew the equation, then the C mark is scored. A C mark is not awarded if a candidate makes two points which contradict each other. Points which are wrong but irrelevant are ignored.
- A marks** A marks are accuracy or answer marks which either depend on an M mark, or which are one of the ways which allow a C mark to be scored. A marks are commonly awarded for final answers to numerical questions. If a final numerical answer, eligible for A marks, is correct, with the correct unit and an acceptable number of significant figures, all the marks for that question are normally awarded. It is very occasionally possible to arrive at a correct answer by an entirely wrong approach. In these rare circumstances, do not award the A marks, but award C marks on their merits. An A mark following an M mark is a dependent mark.
- Brackets ()** around words or units in the mark scheme are intended to indicate wording used to clarify the mark scheme, but the marks do not depend on seeing the words or units in brackets, e.g. 10 (J) means that the mark is scored for 10, regardless of the unit given.
- Underlining** indicates that this must be seen in the answer offered, or something very similar.
- OR/or** indicates alternative answers, any one of which is satisfactory for scoring the mark.
- e.e.o.o.** means "each error or omission".
- o.w.t.t.e.** means "or words to that effect".
- Spelling** Be generous about spelling and use of English. If an answer can be understood to mean what we want, give credit. However, do not allow ambiguities, e.g. spelling which suggests confusion between reflection/refraction/diffraction or thermistor/transistor/transformer.
- Ignore** indicates that something which is not correct or is irrelevant is to be disregarded and does not cause a right plus wrong penalty.
- Not/NOT** indicates that an incorrect answer is not to be disregarded, but cancels another otherwise correct alternative offered by the candidate, i.e. right plus wrong penalty applies.
- cao** correct answer only
- AND** indicates that both answers are required to score the mark.

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0625	32

- ecf meaning "error carried forward" is mainly applicable to numerical questions, but may in particular circumstances be applied in non-numerical questions. This indicates that if a candidate has made an earlier mistake and has carried an incorrect value forward to subsequent stages of working, marks indicated by ecf may be awarded, provided the subsequent working is correct, bearing in mind the earlier mistake. This prevents a candidate being penalised more than once for a particular mistake, but **only** applies to marks annotated ecf.
- Significant Figures** Answers are normally acceptable to any number of significant figures ≥ 2 . Any exceptions to this general rule will be specified in the mark scheme.
- Units** Deduct one mark for each incorrect or missing unit from an answer that would otherwise gain all the marks available for that answer: maximum 1 per question. No deduction is incurred if the unit is missing from the final answer but is shown correctly in the working. Condone wrong use of upper and lower case symbols, e.g. pA for Pa.
- Fractions** Only accept these where specified in the mark scheme.

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0625	32

- 1 (a) (i) (it/comet) travels in a straight line B1
- (ii) area (under graph) OR $s = vt$ in any form OR vt C1
220 000 m OR 220 km A1
- (b) negative acceleration OR deceleration OR (it/the comet) is slowing down B1
acceleration/deceleration (only accept **it** if acc/decel already mentioned)
not constant allow either increasing or decreasing B1
- (c) attempt at gradient OR $(a =) \Delta v / \Delta t$ OR $(0-)12\,000 / 2.0$ OR other correct values for $\Delta v / \Delta t$ C1
 $(-)6000 \text{ m/s}^2$ tolerance 5000 – 7000 m/s^2 A1
- (d) (it/comet) hits surface (of planet) B1
OR stops o.w.t.t.e. B1

[Total: 8]

- 2 (a) $d = m/V$ in any form OR $(V =) m/d$ OR $200/8.4$ C1
 24 cm^3 A1
- (b) (i) density less (than water) OR upthrust \geq weight B1
- (ii) determine any volume of any liquid (V_1) B1
states viable method to submerge wood B1
reads volume (V_2) from previous line and determines volume
of (wood + brass) ($V_2 - V_1$) B1
subtract volume of brass from above (to give volume of wood) B1

[Total: 7]

- 3 (a) (i) (power =) work (done)/time (taken) OR energy (supplied)/time (taken) OR rate of doing
work OR rate of supplying energy B1
- (ii) box 2 (force acting on the object) AND box 5 (distance moved by the object) B1
- (b) (i) multiplies mass of all passengers by h C1
(increase in gpe =) mgh OR uses $12 \times 650 \times 150$ C1
(power = increase in) gpe/time C1
 $1.8 \times 10^4 \text{ W}$ OR 18 kW A1
- (ii) energy to raise the lift OR weight/load/mass of lift OR more weight/load/mass B1

[Total: 7]

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0625	32

- 4 (a) 2 vectors correct direction AND relative length by eye B1
 correct triangle OR rectangle with resultant on correct diagonal B1
 7.2 kN tolerance 7.0 – 7.4 kN B1
- (b) (i) (moment =) force \times distance C1
 (moment = 11 000 \times 1.8 =) 20 kNm A1
- (ii) (moment of weight = 19 000 \times 1.25 =) 24 (kNm) B1
 correct statement based on two moments seen B1

[Total: 7]

- 5 (a) in all parts accept by implication reference to X
 e.g. in (i) accept “it covers a greater range of temperature”
- (i) X covers greater range of temperature OR (goes to) higher temperature OR greater range expressed numerically B1
- (ii) liquid in X expands uniformly (with temperature rise) B1
- (iii) (for the same temperature rise,) the liquid in X expands more B1
- (b) (i) two junctions correctly connected to each other and to meter OR one junction between wires and other junction at connection to meter M1
 temperature difference between junctions A1
 two wires correctly labelled as made of different materials, accept labels metal A & metal B NOT 3 different metals labelled B1
- (ii) junction (in liquid) has low mass / small heat capacity / small size B1
 temperature of junction reacts quickly / quickly reaches temperature of liquid / heat or cools faster B1

[Total: 8]

- 6 (a) suitable particles and fluid, and labelled, in suitable container
 e.g. pollen and water (surface), smoke in air M1
 microscope AND, if smoke used, illumination A1
- (b) movement of particles NOT atoms or molecules B1
 reasonable description of movement
 OR any mention / clear description of movement in different directions
 accept if diagram drawn B1
- (c) collisions between molecules and particles B1
 random movement of molecules OR causes (random) motion of particles B1

[Total: 6]

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0625	32

- 7 (a) pressure high/increased OR molecules/particles close(r/st together) B1
- (b) (i) 1.7 m B1
- (ii) $v = f \lambda$ in any form OR $(f =) v/\lambda$ OR $5/0.025$ C1
200 Hz A1
- (c) three compressions at $23^\circ - 33^\circ$ to wall B1
constant and correct wavelength by eye
only scored if at $8^\circ - 48^\circ$ to wall B1
- (d) (wavelength) greater B1
change of speed correctly related to change of wavelength B1
- [Total: 8]**
- 8 (a) (i) correct O label B1
- (ii) correct I label B1
- (iii) correct F label, allow correctly labelled dot to left of lens B1
- (iv) correct arrows on both rays, anywhere on each ray B1
- (b) $1/n = \sin i/\sin r$ OR $n = \sin i/\sin r$ in any form OR $\sin i/n$ OR $n \sin i$ C1
($r = \sin^{-1}((\sin 35)/1.5) = 22^\circ$)
accept if in diagram A1
emergent ray drawn with $27^\circ \geq r \geq 18^\circ$ B1
- [Total: 7]**
- 9 (a) (i) NAND B1
- (ii) output and one input correctly labelled B1
- (b) rectangle with longitudinal line in middle third, no input or output wire required B1
- (c) (i) temperature (decreases) B1
- (ii) correctly relates change of resistance to change of temperature B1
voltage of mid-point (of potential divider)/left of LED increases OR higher V across
thermistor B1
current flows through/enough V to light LED B1

Page 7	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0625	32

- (d) $1/R_p = 1/R_1 + 1/R_2$ or $(R_p) = R_1R_2/(R_1 + R_2)$ C1
 $(R = 1/(1/4 - 1/6) =) 12 \Omega$ A1

[Total: 9]

- 10 (a) ≥ 3 horizontal lines in gap by eye B1
 ≥ 4 evenly spaced horizontal lines filling $\frac{3}{4}$ of width of gap AND arrows L to R B1

- (b) (i) ammeter deflects / gives a reading OR registers a current B1
 wire cuts the field lines o.w.t.t.e. M1
 e.m.f. / voltage / current induced / produced / generated A1

- (ii) 1 reading / deflection / current increased B1
 2 reading / deflection / current reversed ignore magnitude B1

[Total: 7]

- 11 (a) (i) 800 counts/s B1

- (ii) $\frac{1}{4}$ of (i) B1

- (b) sample 1 γ B1
 sample 2 β NOT γ as extra B1
 sample 3 α NOT extras B1

- (c) α B1

[Total: 6]